arXiv:1407.7160v1 [math.FA] 26 Jul 2014

On extensions of J-skew-symmetric and J-isometric operators.

S.M. Zagorodnyuk

1 Introduction.

Last years an increasing number of papers was devoted to the investigations of operators related to a conjugation in a Hilbert space, see, e.g. [1], [2], [3], [4] and references therein. A conjugation J in a Hilbert space H is an *antilinear* operator on H such that $J^2x = x$, $x \in H$, and $(Jx, Jy)_H = (y, x)_H$, $x, y \in H$. The conjugation J generates the following bilinear form:

$$[x, y]_J := (x, Jy)_H, \qquad x, y \in H.$$

A linear operator A in H is said to be J-symmetric (J-skew-symmetric) if

$$[Ax, y]_J = [x, Ay]_J, \qquad x, y \in D(A), \tag{1}$$

or, respectively,

$$[Ax, y]_J = -[x, Ay]_J, \qquad x, y \in D(A). \tag{2}$$

A linear operator A in H is said to be J-isometric if

$$[Ax, Ay]_J = [x, y]_J, \qquad x, y \in D(A). \tag{3}$$

If $\overline{D(A)} = H$, then conditions (1), (2) and (3) are equivalent to the following conditions:

$$JAJ \subseteq A^*, \tag{4}$$

$$JAJ \subseteq -A^*, \tag{5}$$

and

$$JA^{-1}J \subseteq A^*, \tag{6}$$

respectively. A linear operator A in H is called J-self-adjoint (J-skew-self-adjoint, or J-unitary) if

$$JAJ = A^*, \tag{7}$$

$$JAJ = -A^*, \tag{8}$$

or

$$JA^{-1}J = A^*, \tag{9}$$

respectively.

We shall prove that each densely defined J-skew-symmetric operator (each J-isometric operator with $\overline{D(A)} = \overline{R(A)} = H$) in a Hilbert space H has a J-skew-self-adjoint (respectively J-unitary) extension in a Hilbert space $\widetilde{H} \supseteq H$. We shall follow the ideas of Galindo in [5] with necessary modifications. In particular, Lemma in [5] can not be applied in our case, since its assumptions can never be satisfied with T: $T^2 = I$. In fact, in this case T would be a conjugation in H. Choosing an element $f \in H$ of an orthonormal basis in H which corresponds to T, we would get $(f, Tf) = (f, f) = 1 \neq 0$. Moreover, an exit out of the original space can appear in our case.

We notice that under stronger assumptions on a J-skew-symmetric operator the existence of a J-skew-self-adjoint extension was proved by Kalinina in [6].

Notations. As usual, we denote by $\mathbb{R}, \mathbb{C}, \mathbb{N}, \mathbb{Z}, \mathbb{Z}_+$, the sets of real numbers, complex numbers, positive integers, integers and non-negative integers, respectively. Set $\overline{0, d} = \{0, 1, ..., d\}$, if $d \in \mathbb{N}$; $\overline{0, \infty} = \mathbb{Z}_+$. If H is a Hilbert space then $(\cdot, \cdot)_H$ and $\|\cdot\|_H$ mean the scalar product and the norm in H, respectively. Indices may be omitted in obvious cases. For a linear operator A in H, we denote by $D(A)$ its domain, by $R(A)$ its range, and A^* means the adjoint operator if it exists. If A is invertible then A^{-1} means its inverse. For a set $M \subseteq H$ we denote by $\overline{M}$ the closure of M in the norm of H. By $\operatorname{Lin} M$ we denote the set of all linear combinations of elements of M, and $\operatorname{span} M := \overline{\operatorname{Lin} M}$. By E_H we denote the identity operator in H, i.e. $E_H x = x$, $x \in H$. In obvious cases we may omit the index H. All appearing Hilbert spaces are assumed to be separable.

2 Extensions of J-skew-symmetric and J-isometric operators.

We shall make use of the following lemma.

Lemma 1 *Let H be a Hilbert space with a positive even or infinite dimension, and J be a conjugation on H. Then there exists a subspace M in H such that*

$$M \oplus JM = H.$$

Proof. Let $\{f_n\}_{n=0}^{2d+1}$ be an orthonormal basis in H corresponding to J, i.e. such that $Jf_n = f_n$, $0 \leq n \leq d$; $d \in Z_+ \cup \{+\infty\}$ $(2d + 2 = \dim H)$. Set

$$f^+_{2k,2k+1} = \frac{1}{\sqrt{2}}(f_{2k} + if_{2k+1}), \quad f^-_{2k,2k+1} = \frac{1}{\sqrt{2}}(f_{2k} - if_{2k+1}), \qquad k \in \overline{0, d}.$$

It is easy to see that $\{f^+_{2k,2k+1}, f^-_{2k,2k+1}\}_{k=0}^d$ is an orthonormal basis in H. Set $M := \mathrm{span}\{f^+_{2k,2k+1}\}_{k=0}^d$. It remains to notice that $JM = \mathrm{span}\{f^-_{2k,2k+1}\}_{k=0}^d$. □

Theorem 1 *Let H be a Hilbert space and J be a conjugation on H. Let A be a J-skew-symmetric (J-isometric) operator in H. Suppose that $\overline{D(A)} = H$ (respectively $\overline{D(A)} = \overline{R(A)} = H$). Then there exists a J-skew-self-adjoint (respectively J-unitary) extension of A in a Hilbert space $\widetilde{H} \supseteq H$ (with an extension of J to a conjugation on $\widetilde{H}$).*

Proof. Let A be such an operator as that in the statement of the theorem. The operator A admits the closure which is J-skew-symmetric (respectively J-isometric) (see, e.g. [3, p. 18]). Thus, without loss of generality we shall assume that A is closed. In what follows, in the case of a J-skew-symmetric (J-isometric) A, we shall say about case (a) (respectively case (b)). Set $H_2 = H \oplus H$, and consider the following transformations on H_2:

$$J_2\{x, y\} = \{Jx, Jy\},\ V\{x, y\} = \{y, -x\},\ U\{x, y\} = \{y, x\}, \quad \forall \{x, y\} \in H_2,$$

and $R := UJ_2 = J_2U$, $K := VR$. Observe that R and K are conjugations on H_2. The graph of an arbitrary linear operator C in the Hilbert space H will be denoted by G_C ($\subseteq H_2$). Observe that

$$J_2G_C = G_{JCJ}, \quad RG_C = UG_{JCJ}. \tag{10}$$

If $\overline{D(C)} = H$, then

$$G_{C^*} = H_2 \ominus VG_C. \tag{11}$$

In the case (a) we may write:

$$(\{x, Ax\}, \{JAJy, y\}) = (x, JAJy) + (Ax, y) = 0, \quad \forall x \in D(A), y \in D(JAJ).$$

Then

$$G_A \perp RG_A. \tag{12}$$

In the case (b), we have

$$(\{x, Ax\}, \{JA^{-1}Jy, -y\}) = 0, \quad \forall x \in D(A), y \in D(JA^{-1}J),$$

and therefore

$$G_A \perp KG_A. \tag{13}$$

Set $D = \begin{cases} H_2 \ominus [G_A \oplus RG_A] & \text{in the case (a)} \\ H_2 \ominus [G_A \oplus KG_A] & \text{in the case (b)} \end{cases}$. If $D = \{0\}$ then it means that A is J-skew-self-adjoint (respectively J-unitary), see considerations for

the operator B below. In the opposite case, we have $RD = D$ (respectively $KD = D$).

At first, suppose that D has a positive even or infinite dimension. By Lemma 1 we obtain that there exists a subspace $X \subseteq D$ such that $X \oplus RX = D$ (respectively $X \oplus KX = D$). Since each element of X is orthogonal to $RG_A = VG_{-JAJ}$ ($KG_A = VG_{JA^{-1}J}$), by (11) it follows that

$$X \subseteq G_{-JA^*J} \quad \text{(respectively } X \subseteq G_{J(A^{-1})^*J}\text{)}. \tag{14}$$

Set $G' = G_A \oplus X$. Suppose that $\{0, y\} \in G'$. Then there exist $\{x, Ax\} \in G_A$ such that $\{0, y\} - \{x, Ax\} = \{-x, y - Ax\} \in X$. By (14) we get $y - Ax = JA^*Jx$ (respectively $y - Ax = -J(A^{-1})^*Jx$), and therefore $y = 0$. Thus, G' is a graph G_B of a densely defined linear operator B. Moreover, we have

$$G_B \oplus RG_B = H_2 \quad \text{(respectively } G_B \oplus KG_B = H_2\text{)}.$$

In the case (a) we get

$$UG_B \oplus URG_B = H_2;$$

$$G_{(-B)^*} = H_2 \ominus VG_{-B} = H_2 \ominus UG_B = URG_B = J_2G_B = G_{JBJ}.$$

In the case (b) we get

$$VG_B \oplus VKG_B = H_2;$$

$$G_{B^*} = H_2 \ominus VG_B = VKG_B = -RG_B = G_{JB^{-1}J}.$$

Suppose now that D has a positive odd dimension. In this case we consider a linear operator $\mathcal{A} = A \oplus A$, with $D(\mathcal{A}) = D(A) \oplus D(A)$, in a Hilbert space $\mathcal{H} = H \oplus H$ with a conjugation $\mathcal{J} = J \oplus J$. Observe that $\mathcal{A}$ is a closed $\mathcal{J}$-skew-symmetric ($\mathcal{J}$-isometric) operator with $\overline{D(\mathcal{A})} = \mathcal{H}$ (respectively $\overline{D(\mathcal{A})} = \overline{R(\mathcal{A})} = \mathcal{H}$). Its graph $G_{\mathcal{A}}$ in a Hilbert space $\mathcal{H}_2 = \mathcal{H} \oplus \mathcal{H}$ may be identified with $G_A \oplus G_A$ in $H_2 \oplus H_2$:

$$G_{\mathcal{A}} = \{\{(f, Af), (g, Ag)\},\ f, g \in D(A)\}.$$

Let $\mathcal{R}$, $\mathcal{K}$ be constructed for $\mathcal{A}$ as R and K for A. In the case (a) we see that

$$\mathcal{H}_2 \ominus [G_{\mathcal{A}} \oplus \mathcal{R}G_{\mathcal{A}}] = (H_2 \ominus [G_A \oplus RG_A]) \oplus (H_2 \ominus [G_A \oplus RG_A]),$$

has a positive even dimension. In the case (b), $\mathcal{H}_2 \ominus [G_{\mathcal{A}} \oplus \mathcal{K}G_{\mathcal{A}}]$ has a positive even dimension. Thus, we may apply the above construction with $\mathcal{A}$ instead of A.

□

References

[1] S.R. Garcia, M. Putinar, Complex symmetric operators and applications, Trans. Amer. Math. Soc. **358** (2006), 1285-1315.

[2] S.R. Garcia, M. Putinar, Complex symmetric operators and applications II, Trans. Amer. Math. Soc. **359** (2007), 3913-3931.

[3] S.M. Zagorodnyuk, On a J-polar decomposition of a bounded operator and matrices of J-symmetric and J-skew-symmetric operators, Banach J. Math. Anal. **4**, No. 2 (2010), 11-36.

[4] C.G. Li, T.T. Zhou, Skew symmetry of a class of operators, Banach J. Math. Anal. **8**, no. 1 (2014), 279-294.

[5] A. Galindo, On the existence of J-self-adjoint extensions of J-symmetric operators with adjoint, Communications on pure and applied mathematics **XV** (1962), 423-425.

[6] T.B. Kalinina, *One extension of an operator in a Hilbert space with an anti-unitary transformation*, Functional analysis (Ul'yanovsk) **18** (1982), 63–71 (Russian).

On extensions of J-skew-symmetric and J-isometric operators.

S.M. Zagorodnyuk

In this paper it is proved that each densely defined J-skew-symmetric operator (or each J-isometric operator with $\overline{D(A)} = \overline{R(A)} = H$) in a Hilbert space H has a J-skew-self-adjoint (respectively J-unitary) extension in a Hilbert space $\widetilde{H} \supseteq H$. We follow the ideas of Galindo in [A. Galindo, On the existence of J-self-adjoint extensions of J-symmetric operators with adjoint, Communications on pure and applied mathematics, Vol. XV, 423-425 (1962)] with necessary modifications.